



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Infectious prions do not induce A^β deposition in an in vivo seeding model

Rasmussen, Jay ; Krasemann, Susanne ; Altmeyden, Hermann ; Schwarz, Petra ; Schelle, Juliane ;
Aguzzi, Adriano ; Glatzel, Markus ; Jucker, Mathias

DOI: <https://doi.org/10.1007/s00401-018-1848-5>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-151136>

Journal Article

Accepted Version

Originally published at:

Rasmussen, Jay; Krasemann, Susanne; Altmeyden, Hermann; Schwarz, Petra; Schelle, Juliane; Aguzzi, Adriano; Glatzel, Markus; Jucker, Mathias (2018). Infectious prions do not induce A^β deposition in an in vivo seeding model. *Acta Neuropathologica*, 135(6):965-967.

DOI: <https://doi.org/10.1007/s00401-018-1848-5>

Infectious prions do not induce A β deposition in an *in vivo* seeding model

Jay Rasmussen^{1,2,3,§}, Susanne Krasemann⁴, Hermann Altmeyen⁴, Petra Schwarz⁵, Juliane Schelle^{1,2,3}, Adriano Aguzzi⁵, Markus Glatzel⁴, Mathias Jucker^{1,2,§}

¹German Center for Neurodegenerative Diseases (DZNE), Tübingen, D-72076 Tübingen, Germany; ²Department of Cellular Neurology, Hertie Institute for Clinical Brain Research, University of Tübingen, D-72076 Tübingen, Germany; ³Graduate Training Center of Neuroscience - Cellular & Molecular Neuroscience, University of Tübingen, D-72074 Tübingen, Germany; ⁴Institute of Neuropathology, University Medical Center Hamburg-Eppendorf UKE, D-20246 Hamburg, Germany; ⁵Institute of Neuropathology, University Hospital Zurich, 8091 Zurich, Switzerland.

§ To whom correspondence should be addressed

jdoug.rasmussen@gmail.com; mathias.jucker@uni-tuebingen.de

An increasing number of studies have suggested that certain cases of iatrogenic Creutzfeldt-Jakob disease (iCJD) that harbor significant β -amyloid (A β) pathology are the result of aggregated A β transmission to patients during the same procedure that caused prion disease [2, 4, 7, 8, 11, 13, 17]. The source of iatrogenic contamination has been observed both for human growth hormone infusions and dura mater grafts, arguing against a treatment specific effect. Intriguingly, recent work has also observed suspected A β pathology transmission in post-mortem samples that received growth hormone treatments but did not develop CJD [17]. Yet another study suggested that neurosurgery with A β -contaminated tools can transmit A β pathology and lead to intracerebral hemorrhage [12]. These findings have been debated in the context of whether A β pathology is truly transmissible and whether Alzheimer's disease could subsequently develop.

It is well known that aggregated A β can nucleate the misfolding and aggregation of naïve A β monomers both *in vitro* and *in vivo*, in a process termed seeding [15, 20]. Thus, it appears plausible that exogenous A β seeds could induce pathology in human subjects. However, it has been reported that the cellular prion protein, PrP^C, can bind different A β species, especially oligomers [3, 14]. Furthermore, conflicting reports have suggested that infectious prions (PrP^{Sc}) can exacerbate A β deposition at a time point where plaques are

already present, with both pathologies working synergistically, but also that misfolded A β can actually interfere with prion pathogenesis [16, 18, 19]. Given these findings, it is important to consider that PrP^{Sc} inoculations could influence A β pathology, but the role of PrP^{Sc} in initiating cerebral β -amyloidosis is still unclear. To this end, we inoculated APP transgenic mice on a PrP^C wildtype or heterozygous background with infectious PrP^{Sc} prions to investigate whether PrP^{Sc} alters the onset of A β pathology.

APP23 mice expressing human amyloid precursor protein (APP) with a Swedish mutation (KM670/671NL) were intracerebrally inoculated (hippocampus, bregma: 2.5 mm posterior; +/-2.0 mm lateral; 1.8 mm ventral) with infectious RML-PrP^{Sc} or wildtype (WT) brain extracts (1% w/v) prepared from RML-PrP^{Sc} infected or healthy CD1 mice respectively, using the Precellys system (5500 rpm, 2x20 sec, Bertin Instruments) (Fig. 1a, Supplemental Methods). All PrP^{Sc}-inoculated mice developed terminal prion disease and were sacrificed after 176 days post injection (median). Histological analysis with Hematoxylin & Eosin and SAF84 (PrP^{Sc}, 1:250) revealed that brains displayed typical vacuolation (spongiosis) and PrP-immunoreactive deposits in sick animals, which was not detectable in WT-injected animals of the same age (Fig. 1a). Using an in-house pan-A β antibody (CN6, 1:1000) (Supplemental Methods), no induced A β pathology was detected in the hippocampus of PrP^{Sc} prion-injected animals while, conversely, the injection of minimal A β seeds and a similar incubation period does yield A β deposition as demonstrated previously [20].

Given that the incubation period of the suggested transmission of human A β pathology is 10 to 40 years for both transmission via contaminated dura grafts and growth hormone [2, 4, 8, 10, 11], we hypothesized that the incubation period in our mouse model may not be long enough to detect induced A β pathology caused by PrP^{Sc} prions. Thus, APP23 mice were crossed to a *Prnp*-null line to produce PrP^C heterozygous mice (APP23-PrP^{+/-}), which is known to increase the prion incubation period until terminal sickness [1, 6]. Indeed, APP23-PrP^{+/-} mice injected with PrP^{Sc} prions survived an extra 69 days after intrahippocampal inoculation as described above (median survival: 245 days post injection) before being sacrificed due to prion disease with the expected PrP^{Sc}-positive staining using SAF84 (Fig. 1b). Nevertheless, none of the animals injected with PrP^{Sc} prions presented with detectable seeded A β deposition after staining sections for A β (CN6) (Fig. 1b). In parallel, APP23-PrP^{+/-} mice were also inoculated with brain extracts prepared from aged APP23 A β -laden brains homogenized with the Precellys system as above (10% w/v) followed by a 3000

g centrifugation (5 min). All A β injected animals showed obvious induced A β pathology as expected [15, 20] (Fig. 1b).

The potential of A β pathology transmission under specific circumstances in humans is interesting both from a basic biology and human health perspective [2, 4, 7, 8, 10-13, 17]. However, given that these human studies are observational, there are questions on the mechanism behind the increased A β deposition. We have found that intrahippocampal inoculations of infectious RML prions into APP23 transgenic mice caused prion disease but did not induce A β pathology in the hippocampus after long incubation periods that are sufficient to detect seeded A β pathology caused by nanomolar amounts of A β (one seeding unit) [20]. This argues against a direct cross-seeding effect of PrP^{Sc} prions on A β or an indirect effect of prion disease leading to A β pathology. This conclusion is supported by the recent report that patients who received growth hormone treatments but did not have prion disease still contained significant A β pathology and other instances suggesting A β pathology in prion disease patients is an age-related phenomenon [9, 17].

It is worth noting that, although PrP^{Sc} did not induce A β pathology before terminal prion disease, these inoculations were intracerebral and were meant to provide a model system for dura mater grafting or for contaminated surgical instruments. Instances of A β pathology after peripheral growth hormone treatment in humans with iCJD could be caused by A β seeds in the growth hormone extract traveling to the brain similar to studies in APP transgenic mice [2, 4, 5, 11, 17]. However, it cannot be excluded that a peripheral prion infection indirectly influences A β pathology [18], and thus mechanistically would contrast intracerebral exposure. It is also important to consider that a different PrP^{Sc} prion strain may harbor A β pathology inducing activity.

From our work, we can conclude that misfolded A β introduced during treatment may be responsible for induction of A β pathology in these human cases as opposed to being the by-product of iCJD prion infection. Future work will need to determine whether cases of A β pathology transmission could eventually develop into clinical Alzheimer's disease.

References

- 1 Bueler H, Fischer M, Lang Y, Bluethmann H, Lipp HP, DeArmond SJ, Prusiner SB, Aguet M, Weissmann C (1992) Normal development and behaviour of mice lacking the neuronal cell-surface PrP protein. *Nature* 356: 577-582. doi: 10.1038/356577a0
- 2 Cali I, Cohen ML, Haslam J, Pardi P, Giaccone G, Collins SJ, Kofsky D, Wang H, McLean CA, Brandel JP, et al. (2018) Iatrogenic Creutzfeldt-Jakob disease with Amyloid-beta pathology: an international study. *Acta Neuropathol Commun* 6: 5. doi: 10.1186/s40478-017-0503-z
- 3 Dohler F, Sepulveda-Falla D, Krasemann S, Altmeppen H, Schluter H, Hildebrand D, Zerr I, Matschke J, Glatzel M (2014) High molecular mass assemblies of amyloid-beta oligomers bind prion protein in patients with Alzheimer's disease. *Brain* 137: 873-886. doi: 10.1093/brain/awt375
- 4 Duyckaerts C, Sazdovitch V, Ando K, Seilhean D, Privat N, Yilmaz Z, Peckeu L, Amar E, Comoy E, Maceski A, et al. (2018) Neuropathology of iatrogenic Creutzfeldt-Jakob disease and immunoassay of French cadaver-sourced growth hormone batches suggest possible transmission of tauopathy and long incubation periods for the transmission of Abeta pathology. *Acta Neuropathol* 135: 201-212. doi: 10.1007/s00401-017-1791-x
- 5 Eisele YS, Obermuller U, Heilbronner G, Baumann F, Kaeser SA, Wolburg H, Walker LC, Staufenbiel M, Heikenwalder M, Jucker M (2010) Peripherally applied Abeta-containing inoculates induce cerebral beta-amyloidosis. *Science* 330: 980-982. doi: 10.1126/science.1194516
- 6 Fischer M, Rulicke T, Raeber A, Sailer A, Moser M, Oesch B, Brandner S, Aguzzi A, Weissmann C (1996) Prion protein (PrP) with amino-proximal deletions restoring susceptibility of PrP knockout mice to scrapie. *EMBO J* 15: 1255-1264.
- 7 Frontzek K, Lutz MI, Aguzzi A, Kovacs GG, Budka H (2016) Amyloid-beta pathology and cerebral amyloid angiopathy are frequent in iatrogenic Creutzfeldt-Jakob disease after dural grafting. *Swiss Med Wkly* 146: w14287. doi: 10.4414/smw.2016.14287
- 8 Hamaguchi T, Taniguchi Y, Sakai K, Kitamoto T, Takao M, Murayama S, Iwasaki Y, Yoshida M, Shimizu H, Kakita A, et al. (2016) Significant association of cadaveric dura mater grafting with subpial Abeta deposition and meningeal amyloid angiopathy. *Acta Neuropathol* 132: 313-315. doi: 10.1007/s00401-016-1588-3
- 9 Hainfellner JA, Wanschitz J, Jellinger KA, Liberski PP, Gullotta F, Budka H (1998) Coexistence of Alzheimer-type neuropathology in Creutzfeldt-Jakob disease. *Acta Neuropathol* 96: 116-122.
- 10 Herve D, Porche M, Cabrejo L, Guidoux C, Tournier-Lasserre E, Nicolas G, Adle-Biassette H, Plu I, Chabrier H, Duyckaerts C (2018) Fatal Abeta cerebral amyloid angiopathy 4 decades after a dural graft at the age of 2 years. *Acta Neuropathol*: March 5. doi: 10.1007/s00401-018-1828-9
- 11 Jaunmuktane Z, Mead S, Ellis M, Wadsworth JD, Nicoll AJ, Kenny J, Launchbury F, Linehan J, Richard-Loendt A, Walker AS, et al. (2015) Evidence for human transmission of amyloid-beta pathology and cerebral amyloid angiopathy. *Nature* 525: 247-250. doi: 10.1038/nature15369
- 12 Jaunmuktane Z, Quaegebeur A, Taipa R, Viana-Baptista M, Barbosa R, Koriath C, Scot R, Mead S, Brandner S (2018) Evidence of amyloid-beta cerebral amyloid angiopathy transmission through neurosurgery. *Acta Neuropathol*: Feb 15. doi: 10.1007/s00401-018-1822-2

- 13 Kovacs GG, Lutz MI, Ricken G, Strobel T, Hoftberger R, Preusser M, Regelsberger G, Honigschnabl S, Reiner A, Fischer P, et al (2016) Dura mater is a potential source of Abeta seeds. *Acta Neuropathol* 131: 911-923. doi: 10.1007/s00401-016-1565-x
- 14 Lauren J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM (2009) Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* 457: 1128-1132. doi: 10.1038/nature07761
- 15 Meyer-Luehmann M, Coomaraswamy J, Bolmont T, Kaeser S, Schaefer C, Kilger E, Neuenschwander A, Abramowski D, Frey P, Jaton AL, et al (2006) Exogenous induction of cerebral beta-amyloidogenesis is governed by agent and host. *Science* 313: 1781-1784. doi: 10.1126/science.1131864
- 16 Morales R, Estrada LD, Diaz-Espinoza R, Morales-Scheihing D, Jara MC, Castilla J, Soto C (2010) Molecular cross talk between misfolded proteins in animal models of Alzheimer's and prion diseases. *J Neurosci* 30: 4528-4535. doi: 10.1523/JNEUROSCI.5924-09.2010
- 17 Ritchie DL, Adlard P, Peden AH, Lowrie S, Le Grice M, Burns K, Jackson RJ, Yull H, Keogh MJ, Wei W, et al. (2017) Amyloid-beta accumulation in the CNS in human growth hormone recipients in the UK. *Acta Neuropathol* 134: 221-240. doi: 10.1007/s00401-017-1703-0
- 18 Sarell CJ, Quartermann E, Yip DC, Terry C, Nicoll AJ, Wadsworth JDF, Farrow MA, Walsh DM, Collinge J (2017) Soluble Abeta aggregates can inhibit prion propagation. *Open Biol* 7: pii: 170158. doi: 10.1098/rsob.170158
- 19 Schwarze-Eickner K, Keyvani K, Görtz N, Westaway D, Sachser D, Paulus W (2005) Prion protein (PrPc) promotes beta-amyloid plaque formation. *Neurobiol Aging* 26:1177-82.
- 20 Ye L, Rasmussen J, Kaeser SA, Marzesco AM, Obermuller U, Mahler J, Schelle J, Odenthal J, Kruger C, Fritsch SK, et al. (2017) Abeta seeding potency peaks in the early stages of cerebral beta-amyloidosis. *EMBO Rep* 18: 1536-1544. doi: 10.15252/embr.201744067

Fig. 1. *RML-PrP^{Sc} inoculations into APP23 transgenic mice do not induce A β deposition.* (a) APP23 mice were injected with either RML PrP^{Sc} (n=5; 4 males, 1 female) or a WT control brain extract (n=5; 3 males, 2 females) and monitored until RML-injected animals became sick (median survival: 176 d), at which time point all animals were sacrificed. Representative histological stains for Hematoxylin-Eosin (H&E), PrP^{Sc} (SAF84) and A β (CN6) are presented. None of the RML PrP^{Sc} (0/5) or WT inoculated (0/5) APP23 mice revealed any detectable A β deposits in the hippocampus. H&E, PrP^{Sc} scale bars=100 μ m; A β scale bar=200 μ m. (b) APP23 mice heterozygous for PrP^C (APP23-PrP^{+/-}) were injected with either RML PrP^{Sc} (n=7; 4 males, 3 females), WT control (n=6; 4 males, 2 females) or A β seeding extract (n=6; 3 males, 3 females). Mice were monitored until RML-injected animals became sick, then all animals were sacrificed (median survival: 245 d). Representative histological stainings of PrP^{Sc} (SAF84) and A β (CN6) are presented. None of the RML-PrP^{Sc} (0/7) or WT-inoculated (0/6) APP23-PrP^{+/-} mice revealed any detectable induced A β deposits whereas all A β seed-inoculated (6/6) APP23-PrP^{+/-} mice showed induced A β deposition. PrP^{Sc} scale bar=100 μ m; A β scale bar=200 μ m.